

2003 Health and Preventive Practices Status Report

from Missouri County-level Study



Scott County**

	County Prevalence (%)*	Southeast Region Prevalence (%)*	State Prevalence (%)*
G eneral			
Health Status (Fair or Poor)	24.4 H	23.2 H	16.9
No Health Coverage	16.8	17.5 H	12.8
B ehavioral Risk Factors			
Physical Inactivity	38.7 H	30.8 H	24.0
Current Smoker	24.7	28.1	26.4
Overweight (BMI 25-29.9)	37.3	36.9	35.3
Obese (BMI \geq 30)	29.4	26.9	23.4
D isease and Conditions			
High Blood Pressure	36.5	30.1	28.9
High Cholesterol (35 years or older)	28.7 L	38.2	37.1
Asthma	9.2	12.3	10.8
Diabetes	7.8	7.5	7.2
Activity Limitation	16.2	20.6	17.9
C ancer Screening			
Women			
Never had a Mammogram	14.0	20.5 H	13.4
No Mammogram or Clinical Breast Exam in Last Year:			
40-49 years of age	49.2	58.0	49.4
50-64 years of age	35.6	44.8	35.9
65 and older	50.4	52.9	47.4
Never had a Pap Test:			
18-69 years of age	6.5	3.8	5.1
70 and older	10.9	8.2	6.4
No Pap Test in Last Year (18-29)	39.9	23.5	25.8
No Pap Test in Last 3 Years (30-69)	4.9	15.4	13.8
Men and Women 50 Years or Older			
Blood Stool Test:			
Never had a Blood Stool Test	66.8	63.1 H	57.0
No Blood Stool Test in Last Year	89.0 H	83.7 H	77.5
Sigmoidoscopy/Colonoscopy			
Never had a Sigmoidoscopy or Colonoscopy	62.4 H	59.0 H	49.2
No Sigmoidoscopy or Colonoscopy in Past Five Years	69.3 H	67.7 H	57.0

*Prevalence rates are age adjusted to the 2000 U.S. standard population

■ indicates that the prevalence rate is not significantly above or below state prevalence

H indicates that the prevalence rate is significantly above the state prevalence

L indicates that the prevalence rate is significantly below the state prevalence

Unless otherwise noted respondents are 18 years of age or older

** The best indicators for Scott county were obtained by combining data from Scott and Mississippi counties.